Accepted by ApJ
Preprint typeset using LaTeX style emulateapj v. 11/26/03

arXiv:astro-ph/0401525v1 24 Jan 2004

# THE STATISTICAL PROPERTIES OF GALAXIES CONTAINING ULXS

A. Ptak[1], E. Colbert[1,2]

*Accepted by ApJ*

## ABSTRACT

We present a statistical analysis of the properties of galaxies containing ultraluminous X-ray objects (ULXs). Our primary goal is to establish the fraction of galaxies containing a ULX as a function of ULX luminosity. Our sample is based on *ROSAT* HRI observations of galaxies. We find that $\sim 12\%$ of galaxies contain at least one ULX with $L_X > 10^{39}$ erg s$^{-1}$and $\sim 1\%$ of galaxies contain at least one ULX with $L_X > 10^{40}$ erg s$^{-1}$. These ULX frequencies are lower limits since ROSAT HRI observation would miss absorbed ULXs (i.e., with $N_H \gtrsim 10^{21}$ cm$^{-2}$) and those within $\sim 10$" of the nucleus (due to the positional error circle of the *ROSAT* HRI). The Hubble type distribution of galaxies with a ULX differs significantly from the distribution of types for nearby RC3 galaxies, but does not differ significantly from the galaxy type distribution of galaxies observed by the HRI in general. We find no increase in the mean FIR luminosity or FIR / K band luminosity ratio for galaxies with a ULX relative to galaxies observed by the HRI in general, however this result is also most likely biased by the soft bandpass of the HRI and the relatively low number of high SFR galaxies observed by the HRI with enough sensitivity to detect a ULX.

*Subject headings:*

## 1. INTRODUCTION

*ROSAT* HRI observations showed that ultraluminous X-ray objects (ULXs, a.k.a. IXOs), which were previously detected with *Einstein* (see Fabbiano 1989), are compact X-ray sources, are quite common in the local Universe, and are typically offset from the optical galactic nucleus (Colbert & Mushotzky 1999; Roberts & Warwick 2000). Here we define ULXs as X-ray sources in galaxies that are point-like, extra-nuclear, and have X-ray luminosities in excess of $10^{39}$ ergs s$^{-1}$. There is considerable debate as to nature of these sources (see van der Marel 2003; Miller & Colbert 2003, for reviews), with popular explanations including anisotropic emission from stellar-mass X-ray binaries and intermediate mass black holes. A constraint on any explanation for these sources would be the frequency at which they occur, i.e., what fraction of galaxies harbor a ULX as a function of ULX luminosity. For example, this frequency would have implications for the typical relativistic boosting factor for jet models of ULXs (Koerding, Falcke, & Markoff 2002). In this paper we examine the frequency of ULX occurrence as a function of ULX luminosity based on the HRI ULX catalog published in Colbert & Ptak (2002). While HRI observations suffer from significant biases against the detection of ULXs (as discussed below), we note that the relatively large field of view of the HRI resulted in a large number of galaxies having been observed, and of course large area surveys are crucial for establishing the bright end of a luminosity distribution.

## 2. METHODOLOGY

In addition to Colbert & Ptak (2002), we also made use of a subset of the RC3 galaxy catalog (de Vaucouleurs et al. 1991) where the distance is determined, most often based on Tully (1988). Colbert & Ptak (2002) chose a recessional velocity cutoff of 5000 km s$^{-1}$. In this paper we assume the cosmology $H_0 = 75$ km s$^{-1}$ Mpc$^{-1}$ and $q_0 = 0.5$ (for consistency with Tully 1988), and therefore this velocity corresponds to 66.7 Mpc. This resulted in a galaxy sample containing 9452 galaxies. We next determined which of these galaxies had "coverage" by computing the overlap of the RC3 $D_{25}$ ellipse and the HRI FOV for each observation that was used in Colbert & Ptak (2002). We considered a galaxy as having coverage if $\geq 90\%$ of the galaxy was in the FOV. We found that 766 galaxies had coverage, or $\sim 8\%$ of the RC3 subset.

In order to properly assess the frequency of ULXs, we need to assess how many galaxies were observed by the HRI with sufficient sensitivity to detect a ULX. We computed the flux limit at the position of each RC3 galaxy with coverage by computing the background level in the point source region size. The point source region used by XAssist (Ptak & Griffiths 2003) is a circle with a radius given by $4[\exp(\theta/9)+1]$ arcseconds, where $\theta$ is the off-axis angle of the source in arcminutes. The total number of background counts, b, in this area was then determined using the median background counts per pixel (also computed as a part of the XAssist processing of each field). The $3\sigma$ detection threshold count rate is given by $3\sqrt{b}/t$, where t is the exposure time, and the limiting flux is then $3\sqrt{b} f_{conv} v(\theta)/t$ where $v(\theta)$ gives the vignetting correction and $f_{conv}$ is the on-axis count rate to flux conversion factor. $f_{conv}$ was computed using the FTOOL PIMMS, assuming a power-law slope of 1.7 and the Galactic column density (Dickey & Lockman 1990, using the FTOOL "nh"). We computed the limiting flux in both the 0.5-2.0 keV bandpass $F_{\rm lim,0.5-2.0\ keV}$ and the 2-10 keV bandpass $F_{\rm lim,2-10\ keV}$. We used $F_{\rm lim,2-10\ keV}$ to compute the limiting luminosity since the 2-10 keV bandpass was used in Colbert & Ptak (2002), for consistency with current *Chandra* and *XMM-Newton* surveys.

---

[1] The Johns Hopkins University, Homewood Campus, Baltimore, MD 21218, ptak@pha.jhu.edu

[2] The Catholic University of America, 620 Michigan Ave NE, Washington, DC 20064, colbert@cua.edu

Any ULX could then have been detected in a galaxy when $L_{\rm lim,2-10~keV}$ is $\leq 10^{39}$erg s$^{-1}$, while more generally $L_{\rm lim,2-10~keV}$ is $> 10^{39}$erg s$^{-1}$.

Finally, it is necessary to determine how many ULXs are likely to be background sources. For this purpose we used the logN-logS given in Hasinger (1998), which we note is consistent with a simple power-law $\log N(>S) \sim 100(S_{-14})^{-1.4}$ sources deg$^{-2}$, where $S_{-14}$ is the 0.5-2.0 keV flux in units of $10^{-14}$ ergs cm$^{-2}$ s$^{-1}$. The number of expected background sources is then $n_{xrb} = 100(\frac{F_{\rm lim,0.5-2.0~keV}}{10^{-14}})^{-1.4}\pi r^2$, where r is the major axis radius of each galaxy in degrees. Colbert & Ptak (2002) used $r = D_{25}$ (the RC3 major axis diameter) however here we also consider $r = 0.5D_{25}$. Also note that when $L_{\rm lim,2-10~keV}$ is $\leq 10^{39}$erg s$^{-1}$, the relevant number of background sources is the number of source whose *apparent* luminosity exceeds $10^{39}$erg s$^{-1}$, and since we are assuming a simple power-law logN-logS this simply entails decreasing the total number of background sources expected by the factor $(L_{\rm lim,2-10~keV}/10^{39}{\rm erg~s^{-1}})^{1.4}$.

## 3. RESULTS

### 3.1. *ULX Frequency*

In Table 1 we show the number of galaxies ($N_{L_{lim}\leq L_X}$) with HRI coverage and a limiting 2-10 keV luminosity *less than* or equal to $L_X$ ranging from $10^{39}$erg s$^{-1}$to $10^{41}$erg s$^{-1}$(i.e., the number of galaxies where a ULX as bright as $L_X$ or brighter could have been detected). We also show the number of galaxies with at least one ULX ($N_{wULX}$), and the number of galaxies with at least one ULX whose luminosity exceeds the limiting luminosity ($N_{L_{ULX}\geq L_X}$). This latter quantity, $N_{L_{ULX}\geq L_X}$, is of interest since it represents the number of galaxies with a ULX at least as bright as the list $L_X$, and we then computed the resultant number of galaxies after subtracting the expected number of background sources ($N_{net,L_{ULX}\geq L_X}$, i.e., the number of galaxies where the 1-$\sigma$ lower limit on the number of ULXs is greater than 0.; the lower limit was calculated using the mehology described in Kraft, Burrows, & Nousek (1991), with the background being the expected number of background sources). Finally we list the values $N_{wULX}$, $N_{L_{ULX}\geq L_X}$, and $N_{net,L_{ULX}\geq L_X}$ using both $r = D_{25}$ and $r = 0.5D_{25}$. Note that in the latter case the total number of ULXs in the Colbert & Ptak (2002) sample is reduced from 87 to 49.

In Table 1 shows that a large fraction of ULXs detected at radii larger $0.5D_{25}$ are consistent with background sources, i.e., the *net* number of galaxies with ULXs is often larger when $0.5D_{25}$ is chosen as the limiting radius. Also, in some cases, the number of ULXs detected within the $0.5D_{25}$ radius is above the background level when $r = 0.5D_{25}$ but is not above the background level when $r = D_{25}$ (i.e., since in the latter case the expected number of background sources is a factor of four larger). In Figure 1 we plot the fraction of galaxies harboring a ULX as a function of ULX luminosity ($N_{net,L_{ULX}\geq L_X}/N_{L_{lim}\leq L_X}$) for both $r = D_{25}$ and $r = 0.5D_{25}$. Here the errors were computed according to the methodology described in Kraft, Burrows, & Nousek (1991), with the background taken as the number of galaxies at each luminosity that are consistent with all ULXs being background sources (i.e., $N_{L_{ULX}\geq L_X} - N_{net,L_{ULX}\geq L_X}$). The choice of effective radius for the galaxies does not significantly impact the estimate of ULX frequency once background sources are taken into account. The frequency of ULXs observed by the HRI evidently is $\sim$ 12% and $\sim$ 1% of galaxies harboring a ULX with $L_X >= 10^{39}$erg s$^{-1}$and $L_X >= 10^{40}$erg s$^{-1}$, respectively. We also plot the galaxy fractions for spirals only (i.e., Hubble type in RC3 $\geq 0$). The Hubble type distribution of galaxies with and without a ULX is discussed below, however we note that the fraction of galaxies harboring a ULX is slightly higher in spirals.



FIG. 1.— The frequency of ULX occurrence in galaxies as a function of ULX luminosity. The frequency of ULXs is plotted using effective galactic radii of $D_{25}$ (blue diamonds with dashed error bars) and $0.5D_{25}$ (red crosses with solid error bars). We also plot the frequency observed after limiting the sample to only spirals (with Hubble type >= 0 and radius = $0.5D_{25}$; black filled circles with dotted error bars).

### 3.2. *Hubble Type Distribution*

In Figure 2 we show the Hubble type distribution of the RC3 subsample (discussed in §2 ), the Hubble type distribution of galaxies that had HRI coverage (hereafter HRI galaxies) and the HRI galaxies that contain at least one ULX. The ULX galaxy type distribution is shown both before and after background subtraction, and here we only consider $r < 0.5D_{25}$ ULXs. There is a slight excess of elliptical ULX galaxies relative to the RC3 sample however the larger excess in the HRI galaxy sample suggests that this bias due in part to a bias toward early type galaxies in the HRI observing patterns relative to RC3 galaxies in general. The most conspicuous feature of the ULX galaxy distribution is the excess of T=5-7 galaxies, corresponding to Sc and Scd galaxies. However, in general the ULX galaxy type distributions are not significantly different statistically from the HRI RC3 galaxy type distributions, with the $\chi^2$ tests giving $\chi^2/dof \sim 15/8$, which can only be rejected at the 95% level.

We also investigated the possible connection between star-formation rate (SFR) and ULX frequency by tabulating the far-infrared luminosities (a well-known SFR indicator; Kennicutt 1998) of the galaxy samples dis-

Table 1. Galaxy Number Counts

| | | $r = D_{25}$ | | | $r = 0.5D_{25}$ | | |
|---|---|---|---|---|---|---|---|
| $L_X$ (1) | $N_{L_{lim} \leq L_X}$ (2) | $N_{wULX}$ (3) | $N_{L_{ULX} \geq L_X}$ (4) | $N_{net, L_{ULX} \geq L_X}$ (5) | $N_{wULX}$ (6) | $N_{L_{ULX} \geq L_X}$ (7) | $N_{net, L_{ULX} \geq L_X}$ (8) |
| 1 | 228 | 45 | 45 | 25 (8.7%) | 34 | 34 | 28 (12.3%) |
| 2 | 316 | 49 | 38 | 23 (7.3%) | 37 | 27 | 23 (7.3%) |
| 5 | 446 | 53 | 23 | 13 (2.9%) | 38 | 14 | 14 (3.1%) |
| 10 | 540 | 54 | 11 | 7 (1.3%) | 38 | 6 | 6 (1.1%) |
| 50 | 736 | 54 | 1 | 1 (0.1%) | 38 | 0 | 0 (0.0%) |
| 100 | 762 | 54 | 0 | 0 (0.0%) | 38 | 0 | 0 (0.0%) |

Note. — (1) minimum ULX 2-10 keV luminosity in $10^{39}$erg s$^{-1}$. (2) $N_{L_{lim} \leq L_X}$ = number of galaxies with a limiting luminosity $L_{lim}$ less than or equal to $L_X$. (3) and (6) $N_{wULX}$ = number of galaxies with $L_{lim} \leq L_X$ and a ULX, for $r = D_{25}$ and $r_{0.5D_{25}}$, respectively. (4) and (7) $N_{L_{ULX} \geq L_X}$ = number of galaxies with a $L_{lim} \leq L_X$ and a ULX with luminosity $L_{ULX} \geq L_X$. (5) and (8) $N_{net, L_{ULX} \geq L_X}$ = number of galaxies with a $L_{lim} \leq L_X$ and a net number of ULXs with luminosity $L_{ULX} \geq L_X > 0$ after subtracting the expected number of background sources, with the percentage of galaxies (i.e., $N_{net, L_{ULX} \geq L_X}/N_{L_{lim} \leq L_X}$) given in parentheses.



Fig. 2.— (left) The Hubble type distribution of nearby galaxies in the RC3 catalog (black solid line), the galaxies with HRI coverage (red dashed line), the galaxies with a ULX (blue dot-dashed line) and the galaxies with a ULX after excluding galaxies with at least one "net" ULX (i.e., after subtracting the expected number of background sources; blue dotted line). (right) The Hubble type histograms for galaxies with HRI coverage (red, solid error bars) and with a "net" ULX (blue, dotted error bars) shown with errors computed using Kraft, Burrows, & Nousek (1991). The galaxy types T correspond to: ellipticals have T =-5 to T=-4, lenticulars have T=-3 to T=-1, spirals have T=0 (S0) to T=9, with odd numbers from 1 to 9 representing Sa,b,c,d and even numbers representing Sab,bc,cd,dm. Irregulars have T=10-11 and we assigned 16 to galaxies with T>11. Both plots were created using only ULXs within $r = 0.5D_{25}$.

cussed above. These distributions are shown in Figure 3, where we used NED IRAS fluxes densities at 60 and 100 $\mu m$ to compute $F_{FIR} = 1.26 \times 10^{-23}[2.58 \times 10^{-12} f_{60\mu m} + 10^{-12} f_{100\mu m}]$. The peak of the distributions with a ULX is slightly skewed toward higher FIR (i.e., near $\log L_{FIR} = 43.5$) however the means of the distributions do not differ significantly (all are in the range $\log L_{FIR} = 42.7$-$43.0$). The HRI galaxy FIR luminosity distribution differs from the "net" ULX galaxy FIR luminosity distribution at less than the 90% level based on the $\chi^2$ test. Restricting the sample to spiral galaxies (T >= 0) shifted the means by only $\log L_{FIR} \sim 0.1$, reflecting the predominance of spirals in these samples.

Similarly we plot the distribution of $L_{FIR}/L_K$ where $L_K$ is the K band luminosity computed using the 2mass Extended Source catalog K band magnitude (see Jarrett et al. 2003) and assuming a zero-point flux density of $6.7 \times 10^{-21}$erg cm$^{-2}$ s$^{-1}Hz^{-1}$ (Zombeck 1990)[3]. The K band luminosity is a proxy for stellar mass since the variation in mass-to-light ratio for K band is only a factor of ∼ 2 for various galaxy types (Bell & de Jong 2001). These distributions are shown in Figure 4 and again we do not see any significant differences between galaxies with ULXs and galaxies observed by the HRI in general.

## 4. SUMMARY & CONCLUSIONS

We have assessed the frequency of ULX occurrence in galaxies as a function of ULX luminosity, including proper assessment of the contamination from background sources. We have found that assuming an effective radius of $D_{25}$ tends to result in a significant contamination from

---

[3] see also http://www.ipac.caltech.edu/2mass/releases/allsky/doc/sec6_4a.html



FIG. 3.— The FIR luminosity distribution of nearby galaxies with IRAS detections at 60 $\mu m$ and 100 $\mu m$ (black solid line), the galaxies with HRI coverage (red dashed line), and the galaxies with a ULX before (blue dot-dashed line) and after (blue dotted line) background subtraction. The left panel shows the results without selecting on galaxy type and the right panel shows the results when selecting only spiral galaxies (Hubble type $\geq 0$).



FIG. 4.— The FIR/K band luminosity distributions for nearby galaxies with IRAS detections at 60 $\mu m$ and 100 $\mu m$ and 2mass extended source catalog K band detections (black solid line). Other curves are as in Figure 3. The left panel shows the results without selecting on galaxy type and the right panel shows the results when selecting only spiral galaxies (Hubble type $\geq 0$).

background sources. We found frequencies of $\sim 12\%$ and $\sim 1\%$ for ULX luminosities of at least $10^{39}$erg s$^{-1}$and $10^{40}$erg s$^{-1}$. The Hubble type distribution of galaxies containing at least one ULX is significantly different from those that do not contain a ULX in general, however it is not significantly different from galaxies observed by the HRI. This implies that any bias in the Hubble type distribution of galaxies observed by HRI with ULXs is mostly due to the observing patterns of the HRI. Recently Irwin et al. (2004) reported on an analysis of ULXs in early-type galaxies and found that "very" luminous ULXs (with $L_X > 2 \times 10^{39}$erg s$^{-1}$) are consistent with the expected number of background sources. This result is based in part on *Chandra* data and uses luminosities in the 0.3-10.0 keV bandpass as opposed to our choice of the 2-10 keV band. However, as can be seen in Figure 2b, even when considering all ULXs in our sample (i.e., with $L_X \geq 1 \times 10^{39}$erg s$^{-1}$), the frequency of early-type galaxies containing ULXs is only marginally in excess of zero (with ULXs occurring in globular clusters as in the case of NGC 1399 contributing to the early-type galaxy ULX frequency).

The main caveat with an analysis of the HRI observations is the softness of its bandpass (i.e., 0.1-2.2 keV). Objects absorbed by column densities in excess of $\sim 10^{21}$ cm$^{-2}$ are therefore difficult with the HRI. Since columns of this order are typically observed through the disks of galaxies, it is likely that ULXs are being missed that are close to the nucleus of the galaxies and/or are observed in edge-on disks. *Chandra* and *XMM-Newton* observations will therefore be much more adept at detecting ULXs in these types of environments. This is particularly true for *Chandra* whose much smaller PSF relative to the HRI and *XMM-Newton* allows nuclear ULX to be discerned from AGN at smaller offsets (i.e., 1" compared with 10"). Sipior (2003) finds that 8/41 $\sim 20\%$ of nearby galaxies in a relatively unbiased *Chandra* survey harbor a ULX. This suggests that our HRI survey is missing of order $\sim 40\%$ of ULXs (albeit with large uncertainty due to the size of the Chandra sample). Hornschemeier et al. (2003) reports that the frequency of ULXs with $L_X >= 2 \times 10^{39}$erg s$^{-1}$is $\sim 8\%$ locally and $\sim 36\%$ at z $\sim 0.1$ (Hornschemeier et al. 2003) (suggesting that the frequency of ULXs may evolve). We are similarly finding $\sim 8\%$ (see Table 1) which implies that any biases in our survey are not particularly large for $L_X >= 2 \times 10^{39}$erg s$^{-1}$ULXs, within the errors.

The lack of any obvious connection between FIR luminosity (and hence SFR) or the FIR/K band luminosity ratio distribution and the presence of a ULX suggests that if such a connection exists, then the biases discussed above are particularly acute in star-forming galaxies. This may not be surprising given that high-SFR galaxies tend to contain significant amounts of molecular clouds and dust which would obscure soft X-rays. In addition the relatively small size of our sample may also be impacting this result. For example, only 23 galaxies have $\log L_{FIR}$ >43.5 erg s$^{-1}$and enough HRI sensitivity to detect a ULX, and 4 harbor a ULX (after "background subtraction" and using a galaxy radius of $0.5D_{25}$). This amounts to a larger ULX frequency of $17^{+12}_{-8}\%$, although still statistically consistent with the full sample result. Recent *Chandra* surveys of the point source luminosity functions in galaxies suggest that, particularly at high star-formation rates, the point source luminosity function normalization is correlated with the SFR (Grimm et al. 2003, Colbert et al. 2004), and that the luminosity function is flatter in spirals (Colbert et al. 2004) or (similarly) in higher SFR galaxies (Kilgard et al. 2002). Having both a flatter luminosity function slope and larger normalization at higher SFR strongly implies that higher SFR galaxies (i.e., with SFR greater than several to ten $M_\odot$ yr$^{-1}$ based on Colbert et al. 2004) would tend to harbor *multiple* ULXs. This is also suggested anecdotally by the large numbers of ULXs found in merging galaxies such as the Antennae (Fabbiano, Zezas, & Murray 2001). Accordingly the distribution of $L_{FIR}$ and $L_{FIR}/L_K$ for galaxies with *at least one* ULX may not be significantly impacted by an increase in the total number of ULXs in galaxies with large SFR. However the resolution of these issues will ultimately require more detailed statistical analysis based on *Chandra* and *XMM-Newton* wide-area surveys.

This work was supported by NASA grant NAG 5-11670. We made use of the High Energy Archive and Research Center at NASA/GSFC and the NASA Extragalactic Database. We thanks the anonymous referee for useful comments.

## REFERENCES

Bell, E. & de Jong, R. 2001, ApJ, 550, 212

Colbert, E., Heckman, T., Ptak, A., Strickland, D. & Weaver, K. 2004, ApJ, in press, astro-ph/0305476

Colbert, E. & Ptak, A. 2002, ApJS, 143, 25

Colbert, E. & Mushtozky, R. 1999, ApJ, 519, 89

de Vaucouleurs, G., de Vaucouleurs, A., Corwin, H. G., Jr., Buta, R. J., Paturel, G., & Fouqu, P. 1991, Third Reference Catalog of Bright Galaxies (New York: Springer) (RC3)

Dickey, J. & Lockman, F. 1990, Ann. Rev. Ast. Astr. 28, 215

Fabbiano, G. 1989, ARA&A, 27, 87

Fabbiano, G., Zezas, A. & Murray, S. 2001, ApJ, 554, 1035

Gehrels, N. 1986, ApJ, 303, 336

Grimm, H., Gilfanov, M., Sunyaev, R. 2003, MNRAS, 339, 793

Hasinger, G. 1998, AN, 319, 37

Hornschemeier, A. et al. 2003, ApJ, in press, astro-ph/0308408

Irwin, J., Bregman, J., & Athey, A. 2004, ApJ, in press, astro-ph/0312393)

Jarrett, T., Chester, T., Cutri, R., Schneider, S. E.,& Huchra, J. 2003, AJ, 125, 525

Kennicutt, R. 1998, ARA&A, 36, 189

Kilgard, R. et al. 2002, ApJ, 573, 138

Koerding, E., Falcke, H., & Markoff, S. 2002, A&A, 382, 13

Kraft, R., Burrows, D., & Nousek, J. 1991, ApJ, 374, 344

Miller, M. C. & Colbert, E. 2003, International Journal of Modern Physics D, submitted, astro-ph/0308402

Ptak, A. & Griffiths, R. 2003, Astronomical Data Analysis Software and Systems XII ASP Conference Series, H. E. Payne, R. I. Jedrzejewski, and R. N. Hook, eds., 295, 465

Roberts, T. & Warwick, R. 2000,MNRAS, 304, 52

Sipior, M. 2003, Ph.D. Thesis, Pennsylvania State University

Tully, R. B. 1988, Nearby Galaxies Catalog (Cambridge: Cambridge Univ. Press)

van der Marel, R. 2003, Carnegie Observatories Astrophysics Series, Vol. 1: Coevolution of Black Holes and Galaxies, ed. L. C. Ho (Cambridge: Cambridge Univ. Press), in press, astro-ph/0302101

Zombeck, M. V. 1990, Handbook of Astronomy and Astrophysics, Second Edition (Cambridge, UK: Cambridge University Press)